

Variational approach to the ground state of an impurity in Bose-Einstein condensate

Alexey Novikov¹ and Mikhail Ovchinnikov¹

¹*Department of Chemistry, University of Rochester,
RC Box 270216, Rochester, NY 14627-0216, USA*

(Dated: December 1, 2009)

Abstract

In this paper we consider the effect of self-localization of a quantum impurity in Bose-Einstein condensate. Space correlation function of the impurity is evaluated with the help of the imaginary-time path integral approach. Employing the Feynman's variational method we calculate the impurity correlation function as well as the energy of the system associated with the impurity. The effect of self-localization predicted before within Gross-Pitaevskii approach is recovered by our treatment. The strong coupling regime with negative ground state energy is reached by variational method, and corresponding correlation function is calculated.

I. INTRODUCTION

A theory of atomic size impurity in Bose Einstein fluids has become a subject of intensive research in the last decade. The theoretical development is largely motivated by recent experiments. MIT group [1] has studied the dynamics of impurity atoms in Bose Einstein condensate (BEC) of ultracold atoms in magnetic traps. Another set of experiments studies the microscopic superfluidity of liquid helium by spectroscopic measurements on molecules imbedded in helium droplets [2, 3]. It is generally observed that microscopic impurity particle interacting with Bose liquid/gas behaves as a free particle with effective mass increased by its interaction with Bose fluid. Such dissipationless quantum motion is observed both for translational motion of particles in BEC and for rotations of molecules in superfluid helium.

A question that received attention of theorists has been the structure of the ground state of impurity in BEC, in particular, the possibility of the so called self-localization of an impurity: the appearance of a bound state of an impurity with BEC despite the purely repulsive interaction potential. Such behavior has been investigated by a number of authors using linearized Gross-Pitaevskii equations [4, 5, 6, 7]. This method allows to obtain the non-linear imaginary time Schrödinger equation for the impurity particle. The energy of the particle becomes negative above certain value of coupling strength that depends on the BEC/particle and BEC/BEC interaction potentials. Extension of these results within fully quantum treatment is an open question and is the main subject of present work.

In this paper the matrix element of the reduced density operator of impurity is calculated which allows to compute the energy and the space correlation function of impurity. The system of our interest is the particle interacting with the gas of uncoupled Bogoliubov's excitations. We formulate this matrix element via imaginary time path integrals. The integral over BEC trajectories is explicitly eliminated and the non-Gaussian functional integral over impurity trajectories is left to be calculated. We treat this integral using the variational approach developed by Feynman for the polarons in polar crystals [8]. This method was successfully applied to the problems described by the Fröhlich-type Hamiltonians such as polarons, electron-plasmon interaction [9], nucleon-meson interaction [10], many body fermion problems [11], and many others. Since the Hamiltonian of our model (one particle interacting with uncoupled Bosonic modes) is of the Fröhlich type, it lets us believe that such treatment is successful in our case. We obtain the free energy of an impurity as

a function of coupling constant. The energy is increasing in the weak-coupling regime in agreement with the regular perturbation theory treatments. With further increase of coupling it reaches maximum and decreases becoming negative in the case of strong-coupling. This indicates the existence of the bound state. By computing the correlation function we obtain the localization radius as a function of the coupling constant. We show that the critical localization radius is of the order of magnitude of the inverse critical momentum above which the dissipation of particle motion takes place in real time dynamics.

The Plank constant $\hbar$ and the Boltzmann constant k_B are set to unity throughout the paper.

II. MODEL HAMILTONIAN AND STATEMENT OF A PROBLEM

As it has been mentioned in the Sec. I, we are going to concentrate on the case of dilute Bose gas at the temperature much less than a temperature of condensation. So the gas of weakly interacting Bose particles with mass m can be described as a gas of uncoupled Bogoliubov's excitations [12, 13], i.e. the Hamiltonian of BEC has diagonal form and reads

$$H_B = \sum_{\mathbf{k}} \epsilon(k) \hat{B}_{\mathbf{k}}^+ \hat{B}_{\mathbf{k}} , \quad \epsilon(k) = \sqrt{\frac{k^2}{2m_B} \left(\frac{k^2}{2m_B} + 2m_B c^2 \right)} . \quad (1)$$

Here bosonic operators $\hat{B}_{\mathbf{k}}^+$ and $\hat{B}_{\mathbf{k}}$ create and annihilate the collective excitation with momentum $\mathbf{k}$ and with the Bogoliubov's spectrum $\epsilon(k)$ which has the phonon-like behavior at low momenta, i.e. $\epsilon(k \rightarrow 0) = kc$, where c is the speed of sound.

Next, we consider a quantum particle with mass M interacting with the BEC discussed above. The Hamiltonian of the whole system has the form [14]

$$H = gn + H_B + \sum_{\mathbf{q}} \frac{\mathbf{q}^2}{2M} \hat{a}_{\mathbf{q}}^+ \hat{a}_{\mathbf{q}} + \sum_{\mathbf{q}, \mathbf{k} \neq 0} \gamma_{\mathbf{k}} \left(\hat{a}_{\mathbf{q}-\mathbf{k}}^+ \hat{a}_{\mathbf{q}} \hat{B}_{\mathbf{k}}^+ + \hat{a}_{\mathbf{q}+\mathbf{k}}^+ \hat{a}_{\mathbf{q}} \hat{B}_{\mathbf{k}} \right) , \quad (2)$$

$$\gamma_{\mathbf{k}} = \frac{g}{V} \sqrt{\frac{Nk^2}{2m\epsilon(k)}} .$$

The bosonic operators $\hat{a}_{\mathbf{q}}^+$ and $\hat{a}_{\mathbf{q}}$ in the above expression create and annihilate a single particle with the mass M in the state with momentum $\mathbf{p}$. The last term in Eq. (2) describes the particle-BEC interaction with the coupling constant $\gamma_{\mathbf{q}}$ depending on whole number of particles in Bose gas N in the volume $V = N/n_0$. Also, the coupling strength depends on

the coupling constant, g , defined as a zero Fourier component of impurity/BEC interaction potential. This constant is related to the speed of sound as $c = \sqrt{gn_0/m_B}$. The first term in (2) represents the first order correction to the energy due to the particle-BEC interaction. In the interaction part of the Hamiltonian as well as in the Hamiltonian of the free BEC we neglected the terms responsible for the interaction between the Bogoliubov's excitations. This approximation remains valid if the single impurity alters the surrounding BEC only slightly which is always the case in the macroscopic limit.

In order to describe the statistical properties of the impurity that stays in thermal equilibrium with the BEC environment, we will calculate the space correlation function defined as a matrix element of reduced density operator

$$C(\mathbf{x}, \mathbf{x}') = \frac{1}{Z} \langle \mathbf{x} | \text{Tr}_B e^{-\beta H} | \mathbf{x}' \rangle. \quad (3)$$

Here β is the inverse temperature, the Hamiltonian H is defined by Eq. (2), the trace is performed over all states of BEC and Z is the partition function. Due to the momentum conservation in the whole system the reduced density operator is diagonal in momentum space. Thus

$$C(\mathbf{x} - \mathbf{x}') = \frac{1}{ZV} \sum_{\mathbf{p}} e^{-i\mathbf{p}(\mathbf{x}-\mathbf{x}')} \langle \mathbf{p} | \text{Tr}_B e^{-\beta H} | \mathbf{p} \rangle = \frac{1}{V} \sum_{\mathbf{p}} e^{-i\mathbf{p}(\mathbf{x}-\mathbf{x}')} C_{\mathbf{p}}. \quad (4)$$

So our task is to calculate the correlation function as the reduced density matrix element between two states of the impurity with the momentum $\mathbf{p}$

$$C_{\mathbf{p}} = \rho_{\mathbf{p}}(\beta) = \frac{1}{Z} \text{Tr}_B \langle \mathbf{p} | e^{-\beta H} | \mathbf{p} \rangle = \frac{1}{Z} \text{Tr}_B \langle 0 | \hat{a}_{\mathbf{p}} e^{-\beta H} \hat{a}_{\mathbf{p}}^+ | 0 \rangle \quad (5)$$

The above matrix element can be represented as the imaginary-time coherent-state functional integral [15]

$$\begin{aligned} C_{\mathbf{p}}(\beta) &= \frac{1}{Z} \int D[\{a_{\mathbf{q}}^*(\tau)\}, \{a_{\mathbf{q}}(\tau)\}] \int \prod_{\mathbf{k}} \frac{db_{\mathbf{k}}^* db_{\mathbf{k}}}{\pi} \int D[\{b_{\mathbf{k}}^*(\tau)\}, \{b_{\mathbf{k}}(\tau)\}] a_{\mathbf{p}}(\beta) a_{\mathbf{p}}^*(0) \\ &\times \exp \left[- \sum_{\mathbf{k}} |b_{\mathbf{k}}|^2 + \sum_{\mathbf{k}} b_{\mathbf{k}}^*(\beta) b_{\mathbf{k}}(\beta) - S \right] \end{aligned} \quad (6)$$

with the imaginary-time action of the whole system

$$\begin{aligned} S &= \int_0^\beta d\tau \left[\sum_{\mathbf{q}} (\dot{a}_{\mathbf{q}}(\tau) a_{\mathbf{q}}^*(\tau) + E(\mathbf{q}) a_{\mathbf{q}}^*(\tau) a_{\mathbf{q}}(\tau)) + \sum_{\mathbf{k}} (\dot{b}_{\mathbf{k}}(\tau) b_{\mathbf{k}}^*(\tau) + \epsilon(\mathbf{k}) b_{\mathbf{k}}^*(\tau) b_{\mathbf{k}}(\tau)) \right. \\ &\quad \left. + \sum_{\mathbf{q}, \mathbf{k}} \gamma_{\mathbf{k}} (a_{\mathbf{q}+\mathbf{k}}^*(\tau) a_{\mathbf{q}}(\tau) b_{\mathbf{k}}(\tau) + a_{\mathbf{q}-\mathbf{k}}^*(\tau) a_{\mathbf{q}}(\tau) b_{\mathbf{k}}^*(\tau)) \right]. \end{aligned} \quad (7)$$

The integral (6) must be evaluated with the following boundary conditions for the trajectories

$$b_{\mathbf{k}}(0) = b_{\mathbf{k}} , \quad b_{\mathbf{k}}^*(\beta) = b_{\mathbf{k}}^* , \quad (8)$$

$$a_{\mathbf{q}}(0) = a_{\mathbf{q}}^*(\beta) = 0 . \quad (9)$$

Because of the linear dependence of the interaction part of the action on the $b_{\mathbf{k}}(\tau)$, $b_{\mathbf{k}}^*(\tau)$ trajectories the BEC degrees of freedom can be integrated out immediately using the formula

$$\begin{aligned} & \oint D[\{b_{\mathbf{k}}^*(\tau)\}, \{b_{\mathbf{k}}(\tau)\}] \exp \left[- \sum_{\mathbf{k}} \int_0^\beta d\tau \left(\epsilon(\mathbf{k}) b_{\mathbf{k}}^*(\tau) b_{\mathbf{k}}(\tau) - j_{\mathbf{k}}(\tau) b_{\mathbf{k}}(\tau) - j_{\mathbf{k}}^*(\tau) b_{\mathbf{k}}^*(\tau) \right) \right] \\ &= Z_B \exp \left(\sum_{\mathbf{k}} \int_0^\beta d\tau \int_0^\beta d\tau' \Gamma_{\mathbf{k}}(\tau - \tau') j_{\mathbf{k}}(\tau) j_{\mathbf{k}}^*(\tau') \right) . \end{aligned} \quad (10)$$

The function $\Gamma_{\mathbf{k}}(\tau)$ represents the imaginary-time propagator of free Bose field with the spectrum $\epsilon(\mathbf{k})$

$$\Gamma_{\mathbf{k}}(\tau) = (\Theta(\tau) + n_{\mathbf{k}}) e^{-\tau \epsilon(\mathbf{k})} , \quad n_{\mathbf{k}} = \frac{1}{e^{\beta \epsilon(\mathbf{k})} - 1} , \quad (11)$$

where function $\Theta(\tau)$ denotes the Heaviside step function, and $Z_B = \prod_{\mathbf{k}} n_{\mathbf{k}} e^{\beta \epsilon(\mathbf{k})}$ is its partition function.

Thus, after tracing out the BEC we obtain the functional integral representation of the reduced density matrix of the relevant particle that includes the integration over the particle trajectories only

$$\rho_{\mathbf{p}}(\beta) = \frac{Z_B}{Z} \int D[\{a_{\mathbf{q}}^*(\tau)\}, \{a_{\mathbf{q}}(\tau)\}] a_{\mathbf{p}}(\beta) a_{\mathbf{p}}^*(0) \exp [-S_P + S_I] \quad (12)$$

Here we defined the free imaginary-time action of relevant particle

$$S_P = \int_0^\beta d\tau \sum_{\mathbf{q}} (\dot{a}_{\mathbf{q}}(\tau) a_{\mathbf{q}}^*(\tau) + E(\mathbf{q}) a_{\mathbf{q}}^*(\tau) a_{\mathbf{q}}(\tau)) \quad (13)$$

The integral (12) is of the non-Gaussian type and its non-Gaussian part S_I is due to the impurity-BEC coupling and reads

$$S_I = \int_0^\beta d\tau \int_0^\beta d\tau' \sum_{\mathbf{q}, \mathbf{q}', \mathbf{k}} \gamma_{\mathbf{k}}^2 a_{\mathbf{q}+\mathbf{k}}^*(\tau) a_{\mathbf{q}}(\tau) \Gamma_{\mathbf{k}}(\tau - \tau') a_{\mathbf{q}'-\mathbf{k}}^*(\tau') a_{\mathbf{q}'}(\tau') . \quad (14)$$

Having the action of the impurity, we can investigate its classical dynamics described by corresponding Euler's equations of motion. For this purpose let us write the action

in the coordinate representation by changing the variables of the functional integration as $a_{\mathbf{k}}(\tau) = \frac{1}{\sqrt{V}} \int d\mathbf{r} e^{-i\mathbf{k}\mathbf{r}} \psi(\mathbf{r}, \tau)$. Now the action of the impurity surrounded by BEC has the form

$$S_{imp} = S_P - S_I = \int_0^\beta \int d\mathbf{r} \psi^*(\mathbf{r}, \tau) \left(\frac{\partial}{\partial \tau} - \frac{\Delta}{2M} \right) \psi(\mathbf{r}, \tau) - \int_0^\beta d\tau \int_0^\beta d\tau' \int d\mathbf{r} \int d\mathbf{r}' \psi^*(\mathbf{r}, \tau) \psi(\mathbf{r}, \tau) \Lambda(\mathbf{r} - \mathbf{r}', \tau - \tau') \psi^*(\mathbf{r}', \tau') \psi(\mathbf{r}', \tau') , \quad (15)$$

$$\Lambda(\mathbf{r} - \mathbf{r}', \tau - \tau') = \frac{V}{(2\pi)^3} \int d\mathbf{k} \gamma_{\mathbf{k}}^2 \Gamma_{\mathbf{k}}(\tau - \tau') e^{i\mathbf{k}\mathbf{r}} \quad (16)$$

Varying the above action one gets the following equations of motion

$$\begin{aligned} \dot{\psi}(\mathbf{r}, \tau) &= \frac{\Delta}{2M} \psi(\mathbf{r}, \tau) + \int d\tau' \int d\mathbf{r}' \psi(\mathbf{r}, \tau) \tilde{\Lambda}(\mathbf{r} - \mathbf{r}', \tau - \tau') \psi^*(\mathbf{r}', \tau') \psi(\mathbf{r}', \tau') , \\ -\dot{\psi}^*(\mathbf{r}, \tau) &= \frac{\Delta}{2M} \psi^*(\mathbf{r}, \tau) + \int d\tau' \int d\mathbf{r}' \psi^*(\mathbf{r}, \tau) \tilde{\Lambda}(\mathbf{r} - \mathbf{r}', \tau - \tau') \psi^*(\mathbf{r}', \tau') \psi(\mathbf{r}', \tau') , \end{aligned} \quad (17)$$

with symmetrized kernel

$$\tilde{\Lambda}(\mathbf{r} - \mathbf{r}', \tau - \tau') = \Lambda(\mathbf{r} - \mathbf{r}', \tau - \tau') + \Lambda(\mathbf{r}' - \mathbf{r}, \tau' - \tau) .$$

We will seek the stationary solutions in the form $\psi(\mathbf{r}, \tau) = \psi_s(\mathbf{r}) e^{-\tau E}$ and $\psi^*(\mathbf{r}, \tau) = \psi_s^*(\mathbf{r}) e^{\tau E}$. So the stationary equation for the impurity wave function is

$$E\psi_s(\mathbf{r}) + \frac{\Delta}{2M} \psi_s(\mathbf{r}) + \int d\mathbf{r}' \psi_s(\mathbf{r}) \psi_s^*(\mathbf{r}') \psi_s(\mathbf{r}') \Lambda_s(\mathbf{r} - \mathbf{r}') = 0 , \quad (18)$$

where

$$\Lambda_s(\mathbf{r} - \mathbf{r}') = \int d\tau' \tilde{\Lambda}(\mathbf{r} - \mathbf{r}', \tau - \tau') . \quad (19)$$

Using the definition of the BEC-propagator Eq. (11) together with the Eq. (16) one obtains the non-linear kernel Λ_s , given by

$$\Lambda_s(\mathbf{r}) = \frac{m_B n_0 g^2}{\pi} \frac{e^{-2m_B c r}}{r} . \quad (20)$$

The equations (18) with the kernel (20) represent the particle self-interacting through the screened Coulomb potential. The same equations for the impurity in weak-coupled BEC can also be obtained directly with the help of linearized Gross-Pitaevskii equations with more restricting initial assumptions [5] such as Hartree approximation for the impurity/BEC wave

function and using the real-valued BEC wave function. In Ref. [5] authors investigated the lowest energy solution of the above equations. They found that the equations (18) have the localized solution with negative energy in the strong coupling regime, i.e. when the impurity/BEC interaction is strong enough to compensate the high kinetic energy of the localized state of the impurity.

The aim of this paper is to construct fully quantum description of the model system based on the calculation of the imaginary-time quantum propagator. The quantum formulation of the problem without the restriction of a mean-field approximation will be given in the following section.

III. PATH INTEGRAL FORMULATION OF THE CORRELATION FUNCTION

Our goal is to describe the effect of self-localization from purely quantum point of view. For this purpose let us return to the functional integral for the correlation function with the full impurity-plus-BEC action of the form (6). Our aim is to represent the correlation function as a single Feynman's path integral over the impurity trajectories. As in the previous section we will use the coordinate representation for impurity trajectories. So the correlation function is given by

$$C(\mathbf{x} - \mathbf{x}', \beta) = \frac{1}{Z} \oint D[\{b_{\mathbf{k}}^*(\tau)\}, \{b_{\mathbf{k}}(\tau)\}] \int D[\{\psi^*(\mathbf{r}, \tau)\}, \{\psi(\mathbf{r}, \tau)\}] \psi(\mathbf{x}', \beta) \psi^*(\mathbf{x}, 0) e^{-S} \quad (21)$$

where the action is

$$\begin{aligned} S = & \int_0^\beta d\tau \sum_{\mathbf{k}} (\dot{b}_{\mathbf{k}}(\tau) b_{\mathbf{k}}^*(\tau) + \epsilon(\mathbf{k}) b_{\mathbf{k}}^*(\tau) b_{\mathbf{k}}(\tau)) + \int_0^\beta d\tau \int d\mathbf{r} \left(\psi^*(\mathbf{r}, \tau) \dot{\psi}(\mathbf{r}, \tau) - \psi^*(\mathbf{r}, \tau) \frac{\Delta}{2M} \psi(\mathbf{r}, \tau) \right) \\ & \times \int_0^\beta d\tau \int d\mathbf{r} \psi^*(\mathbf{r}, \tau) \psi(\mathbf{r}, \tau) \sum_{\mathbf{k}} \gamma_{\mathbf{k}} (b_{\mathbf{k}}(\tau) e^{i\mathbf{k}\mathbf{r}} + b_{\mathbf{k}}^*(\tau) e^{-i\mathbf{k}\mathbf{r}}) . \end{aligned} \quad (22)$$

Let us introduce auxiliary external sources into the action and define the following functional

$$\begin{aligned} F[j^*(\mathbf{r}, \tau), j(\mathbf{r}, \tau)] = & \int D[\{\psi^*(\mathbf{r}, \tau)\}, \{\psi(\mathbf{r}, \tau)\}] \\ & \times \exp \left[- \int_0^\beta d\tau \int d\mathbf{r} \left(\psi^*(\mathbf{r}, \tau) \hat{K}(\mathbf{r}, \tau) \psi(\mathbf{r}, \tau) - j(\mathbf{r}, \tau) \psi(\mathbf{r}, \tau) - j^*(\mathbf{r}, \tau) \psi^*(\mathbf{r}, \tau) \right) \right] , \end{aligned} \quad (23)$$

$$\hat{K}(\mathbf{r}, \tau) = \frac{\partial}{\partial \tau} - \frac{\Delta}{2M} + \sum_{\mathbf{k}} \gamma_{\mathbf{k}} (b_{\mathbf{k}}(\tau) e^{i\mathbf{k}\mathbf{r}} + b_{\mathbf{k}}^*(\tau) e^{-i\mathbf{k}\mathbf{r}}) = \frac{\partial}{\partial \tau} + \hat{\mathcal{H}}(\mathbf{r}, \tau) . \quad (24)$$

The functional integral in the right side of Eq. (23) has to be evaluated with the boundary conditions $\psi(\mathbf{r}, 0) = \psi^*(\mathbf{r}, \beta) = 0$. Since the integral in (23) is Gaussian, it can be done by the stationary phase method. Thus for the stationary trajectories one finds

$$\begin{aligned} \dot{\psi}(\mathbf{r}, \tau) - \hat{\mathcal{H}}(\mathbf{r}, \tau)\psi(\mathbf{r}, \tau) - j^*(\mathbf{r}, \tau) &= 0, \\ -\dot{\psi}^*(\mathbf{r}, \tau) - \hat{\mathcal{H}}(\mathbf{r}, \tau)\psi^*(\mathbf{r}, \tau) - j(\mathbf{r}, \tau) &= 0. \end{aligned} \quad (25)$$

The formal solution of the above equations can be written in the following form

$$\psi_s(\mathbf{r}, \tau) = \int_0^\tau d\tau' \exp\left(\int_{\tau'}^\tau \hat{\mathcal{H}}(\mathbf{r}, s) ds\right) j^*(\mathbf{r}, \tau'). \quad (26)$$

Finally, substituting this solution into the integrand in (23), for the functional F we have

$$\begin{aligned} F[j^*(\mathbf{r}, \tau), j(\mathbf{r}, \tau)] &= \exp\left(\int_0^\beta d\tau \int d\mathbf{r} j(\mathbf{r}, \tau) \psi_s(\mathbf{r}, \tau)\right) \\ &= \exp\left(\int_0^\beta d\tau \int_0^\beta d\tau' \Theta(\tau - \tau') \langle j(\mathbf{r}, \tau) | e^{-\int_{\tau'}^\tau \hat{\mathcal{H}}(\mathbf{r}, s) ds} | j^*(\mathbf{r}, \tau') \rangle\right) \end{aligned} \quad (27)$$

Now we can note that the correlation function can be written with the help of the generating functional, namely

$$C(\mathbf{x} - \mathbf{x}', \beta) = \left\langle \frac{\delta^2}{\delta j(\mathbf{x}', \beta) \delta j^*(\mathbf{x}, 0)} F[j^*(\mathbf{r}, \tau), j(\mathbf{r}, \tau)] \right\rangle \Bigg|_{j=j^*=0}, \quad (28)$$

where the averaging is performed as the integration over BEC trajectories with the free action of BEC as

$$\langle \dots \rangle = \frac{1}{Z} \oint D[\{b_{\mathbf{k}}^*(\tau)\}, \{b_{\mathbf{k}}(\tau)\}] \dots \exp\left[\int_0^\beta d\tau \sum_{\mathbf{k}} (\dot{b}_{\mathbf{k}}(\tau) b_{\mathbf{k}}^*(\tau) + \epsilon(\mathbf{k}) b_{\mathbf{k}}^*(\tau) b_{\mathbf{k}}(\tau))\right]. \quad (29)$$

Calculating the functional derivative in Eq. (28) and writing one particle propagator $\langle j(\mathbf{r}, \tau) | e^{-\int_{\tau'}^\tau \hat{\mathcal{H}}(\mathbf{r}, s) ds} | j^*(\mathbf{r}, \tau') \rangle$ as Feynman's path integral, for the correlation function we have

$$C(\mathbf{x} - \mathbf{x}', \beta) = \left\langle \int_{\mathbf{x}(0)=\mathbf{x}}^{\mathbf{x}(\beta)=\mathbf{x}'} D[\mathbf{x}(\tau)] e^{-S_p} \right\rangle, \quad (30)$$

where the single particle action depends on the coordinate trajectory $\mathbf{x}(\tau)$

$$S_p = \int_0^\beta d\tau \left[\frac{M \dot{\mathbf{x}}^2}{2} + \sum_{\mathbf{k}} \gamma_{\mathbf{k}} (b_{\mathbf{k}}(\tau) e^{i\mathbf{k}\mathbf{x}(\tau)} + b_{\mathbf{k}}^*(\tau) e^{-i\mathbf{k}\mathbf{x}(\tau)}) \right] \quad (31)$$

Now using the formula (10) we can eliminate the integration over BEC trajectories from the equation (30) and get the correlation function as a single Feynman's path integral for the impurity

$$C(\mathbf{x} - \mathbf{x}', \beta) = \int_{\mathbf{x}(0)=\mathbf{x}}^{\mathbf{x}(\beta)=\mathbf{x}'} D[\mathbf{x}(\tau)] e^{-S_R} , \quad (32)$$

where the impurity action reads

$$S_R = \int_0^\beta d\tau \frac{M \dot{\mathbf{x}}^2}{2} - \int_0^\beta d\tau \int_0^\beta d\tau' \sum_{\mathbf{k}} \gamma_{\mathbf{k}}^2 \Gamma_{\mathbf{k}}(\tau - \tau') e^{i\mathbf{k}(\mathbf{x}(\tau) - \mathbf{x}(\tau'))} . \quad (33)$$

This form of the single particle functional integral is considered in the following section.

IV. FEYNMAN'S VARIATIONAL APPROACH

The imaginary-time correlation function in the form of path integral (32) provides the exact description of the statistics of the impurity surrounded by the degenerate BEC at low temperature. In this section we will calculate this integral approximately using Feynman's variational approach to the polaron problem [8]. The idea of the original method is to replace the real action of the impurity (33) with the trial action of the form

$$S_T = \int_0^\beta d\tau \frac{M \dot{\mathbf{x}}^2}{2} + \int_0^\beta d\tau \int_0^\beta d\tau' Q(\tau - \tau') (\mathbf{x}(\tau) - \mathbf{x}(\tau'))^2 , \quad (34)$$

where

$$Q(\tau - \tau') = Q_0 (\Theta(\tau - \tau') + n_\omega) e^{-\omega(\tau - \tau')} , \quad n_\omega = \frac{1}{e^{\omega\beta} - 1} . \quad (35)$$

It is also useful to write the trial action in the extended form as

$$e^{-S_T} \sim \oint D[\mathbf{y}(\tau)] e^{-S_{ET}} , \quad (36)$$

$$S_{ET} = \int_0^\beta d\tau \left[\frac{M \dot{\mathbf{x}}^2(\tau)}{2} + \frac{m \dot{\mathbf{y}}^2(\tau)}{2} + \frac{m\omega^2 (\mathbf{y}(\tau) - \mathbf{x}(\tau))^2}{2} \right] . \quad (37)$$

Performing the integration over trajectories $\mathbf{y}(\tau)$ one recovers the trial action in the form of Eq. (34) with $Q_0 = m\omega^3/4$. Thus in this variational treatment we replace the interaction with the original BEC environment by the interaction with single trial particle with the mass m .

The trial action of the form (34) is assumed to be the zeroth order contribution while the difference between real action (33) and trial one has to be considered as a perturbation. Thus one has to write the following expansion of the correlation function

$$C_T(\mathbf{x} - \mathbf{x}', \beta) = \frac{\langle 1 + S_T - S_R \rangle}{\langle 1 + S_T - S_R \rangle_0}. \quad (38)$$

where we have defined two kinds of averages

$$\langle \dots \rangle = \int_{\mathbf{x}(0)=\mathbf{x}}^{\mathbf{x}(\beta)=\mathbf{x}'} D[\mathbf{x}(\tau)] \dots e^{-S_T}, \quad \langle \dots \rangle_0 = \int_{\mathbf{x}(0)=0}^{\mathbf{x}(\beta)=0} D[\mathbf{x}(\tau)] \dots e^{-S_T}. \quad (39)$$

The trial correlation function C_T in Eq. (38) still depends on two variational parameters Q_0 and ω (or m and ω). Thus, in accordance with the principle of minimal sensitivity [16] one has to minimize this expansion with respect to the variational parameters, and the extremum point will give the best variational approximation for the path integral (32). In order to proceed with the expansion (38) we will have to calculate four functions, namely

$$\begin{aligned} \Sigma_1 &= \int_0^\beta d\tau \int_0^\beta d\tau' Q(\tau - \tau') \langle (\mathbf{x}(\tau) - \mathbf{x}(\tau'))^2 \rangle, \\ \Sigma_2 &= \int_0^\beta d\tau \int_0^\beta d\tau' \sum_{\mathbf{k}} \Gamma_{\mathbf{k}}(\tau - \tau') \langle \exp [i\mathbf{k}(\mathbf{x}(\tau) - \mathbf{x}(\tau'))] \rangle, \end{aligned} \quad (40)$$

and

$$\begin{aligned} \Sigma_1^0 &= \int_0^\beta d\tau \int_0^\beta d\tau' Q(\tau - \tau') \langle (\mathbf{x}(\tau) - \mathbf{x}(\tau'))^2 \rangle_0, \\ \Sigma_2^0 &= \int_0^\beta d\tau \int_0^\beta d\tau' \sum_{\mathbf{k}} \Gamma_{\mathbf{k}}(\tau - \tau') \langle \exp [i\mathbf{k}(\mathbf{x}(\tau) - \mathbf{x}(\tau'))] \rangle_0, \end{aligned} \quad (41)$$

The expansion (38) in terms of functions Σ becomes

$$C_T = \frac{\langle 1 \rangle + \Sigma_1 + \Sigma_2}{\langle 1 \rangle_0 + \Sigma_1^0 + \Sigma_2^0} \quad (42)$$

In order to calculate the average $\langle (\mathbf{x}(\tau) - \mathbf{x}(\tau'))^2 \rangle$ and $\langle \exp [i\mathbf{k}(\mathbf{x}(\tau) - \mathbf{x}(\tau'))] \rangle$ that enters the functions Σ_1 and Σ_2 , respectively, one can use the generating functional of the form

$$W[\mathbf{j}(\tau)] = \oint D[\mathbf{y}(\tau)] \int_{\mathbf{x}(0)=\mathbf{x}}^{\mathbf{x}(\beta)=\mathbf{x}'} D[\mathbf{x}(\tau)] \exp \left(-S_{ET} + \int_0^\beta ds \mathbf{x}(s) \mathbf{j}(s) \right) \quad (43)$$

So the second average $\langle \exp [i\mathbf{k}(\mathbf{x}(\tau) - \mathbf{x}(\tau'))] \rangle$ can be evaluated by setting $\mathbf{j}(s) = i\mathbf{k}(\delta(s - \tau) - \delta(s - \tau'))$ while the first one is obtained by differentiating the second one twice over $\mathbf{k}$

at $\mathbf{k} = 0$. The functional integral in (43) can be evaluated by introducing the new variables $\mathbf{q}(\tau) = \mathbf{x}(\tau) - \mathbf{y}(\tau)$ and $\mathbf{r}(\tau) = (M\mathbf{x}(\tau) + m\mathbf{y}(\tau))/(M + m)$. Substituting this replacement into (36) one gets the action S_{ET} in the form

$$S_{ET} = \int_0^\beta d\tau \left[\frac{(M+m)\dot{\mathbf{r}}^2(\tau)}{2} + \frac{\mu\dot{\mathbf{q}}^2(\tau)}{2} + \frac{m\omega^2\mathbf{q}^2(\tau)}{2} - \mathbf{j}(\tau)\mathbf{r}(\tau) - \frac{\mu}{M}\mathbf{j}(\tau)\mathbf{q}(\tau) \right], \quad (44)$$

$$\mu = \frac{mM}{m+M}.$$

The above action now describes harmonic oscillator and free particle in presence of the external source and the corresponding functional integral can be easily evaluated. Here we will write down the result of the integration in (43)

$$W[\mathbf{j}(\tau)](\mathbf{x}, \mathbf{x}') = W_0 \exp \left\{ -(\mathbf{x} - \mathbf{x}')^2 \left[\frac{M\mu}{2\beta m} + \frac{\Omega\mu}{4} \coth \frac{\Omega\beta}{2} \right] \right. \quad (45)$$

$$+ \int_0^\beta d\tau \mathbf{j}(\tau) \left[(\mathbf{x}' - \mathbf{x}) \frac{\mu}{m} \frac{\tau}{\beta} + \frac{\mu}{m} \mathbf{x} + \frac{\mu}{2M} (\mathbf{x} + \mathbf{x}') + \frac{\mu}{2M} \frac{\sinh \Omega\tau - \sinh \Omega(\beta - \tau)}{\sinh \Omega\beta} (\mathbf{x}' - \mathbf{x}) \right]$$

$$+ \int_0^\beta d\tau \int_0^\beta d\tau' \mathbf{j}(\tau) \mathbf{j}(\tau') \left[\frac{\tau'}{2\beta} \frac{\beta - \tau}{M + m} \theta(\tau - \tau') + \frac{\tau}{2\beta} \frac{\beta - \tau'}{M + m} \theta(\tau' - \tau) \right.$$

$$+ \frac{\mu}{4\Omega M^2} \frac{1}{\tanh \frac{\Omega\beta}{2}} \left(\frac{\sinh \Omega(\beta - \tau) + \sinh \Omega\tau + \sinh \Omega(\beta - \tau') + \sinh \Omega\tau'}{\sinh \Omega\beta} - 1 \right)$$

$$\left. + \frac{\mu}{4\Omega M^2} \frac{\cosh \Omega(\beta/2 - |\tau - \tau'|)}{\sinh \frac{\Omega\beta}{2}} \right] \Bigg\}, \quad (46)$$

where we have introduced the frequency $\Omega = \omega\sqrt{m/\mu}$. The prefactor W_0 is the constant coming from the integration over the deviations around the stationary trajectories, so it is independent on the end-points $\mathbf{x}$ and $\mathbf{x}'$.

So for the second average $\langle \exp [i\mathbf{k}(\mathbf{x}(\tau) - \mathbf{x}(\tau')))] \rangle$ one gets

$$\langle \exp [i\mathbf{k}(\mathbf{x}(\tau) - \mathbf{x}(\tau')))] \rangle = W_0 \exp \left\{ -(\mathbf{x} - \mathbf{x}')^2 \left[\frac{M}{2\beta} \frac{\omega^2}{\Omega^2} + \frac{\Omega\mu}{4} \coth \frac{\Omega\beta}{2} \right] \right. \quad (47)$$

$$+ i\mathbf{k}(\mathbf{x}' - \mathbf{x}) \left[\frac{\omega^2}{\Omega^2} \frac{\tau - \tau'}{\beta} + \frac{1}{2} \frac{\Omega^2 - \omega^2}{\Omega^2} \frac{\sinh \Omega\tau - \sinh \Omega(\beta - \tau) - \sinh \Omega\tau' + \sinh \Omega(\beta - \tau')}{\sinh \Omega\beta} \right]$$

$$\left. - \mathbf{k}^2 \left[\frac{1}{2M} \frac{\omega^2}{\Omega^2} |\tau - \tau'| \left(1 - \frac{|\tau - \tau'|}{\beta} \right) + \frac{1}{2M\Omega} \frac{\Omega^2 - \omega^2}{\Omega^2} \frac{\cosh \frac{\Omega\beta}{2} - \cosh \Omega(\beta/2 - |\tau - \tau'|)}{\sinh \frac{\Omega\beta}{2}} \right] \right\}$$

Below we will be always interested in low temperature limit $\beta \rightarrow \infty$. Calculating the first average $\langle (\mathbf{x}(\tau) - \mathbf{x}(\tau'))^2 \rangle$ with the help of the above expression, for the function Σ_1 one gets

$$\Sigma_1 \Big|_{\beta \rightarrow \infty} = W_0 \left[\frac{\Omega^2 + \omega^2}{2\omega\Omega^3} (\mathbf{x} - \mathbf{x}')^2 + \frac{3\beta}{M\omega\Omega} \right] \exp \left[-\frac{\mu\Omega}{4} (\mathbf{x} - \mathbf{x}')^2 \right] \quad (48)$$

The calculation of Σ_2 can not be done in closed form and requires some numerical calculations that will be performed in the next sections.

V. ENERGY OF THE IMPURITY

In this section we will investigate the ground state energy of the impurity as a function of the impurity-BEC coupling strength. Let us define the free energy of the system E as

$$e^{-\beta E} = \int_{\mathbf{x}(0)=0}^{\mathbf{x}(\beta)=0} D[\mathbf{x}(\tau)] e^{-S_R} = \langle e^{S_T - S_R} \rangle_0. \quad (49)$$

In the zero temperature limit free energy coincides with the ordinary energy and shows the difference between coupled and uncoupled impurity/BEC system.

The expression (49) up to first cumulant reads

$$e^{-\beta E(\Omega, \omega)} = \langle 1 \rangle_0 \exp \frac{\langle S_T - S_R \rangle_0}{\langle 1 \rangle_0}, \quad E(\Omega, \omega) = E_0 + E_1 + E_2. \quad (50)$$

The zero energy contribution E_0 is

$$e^{-\beta E_0} = \int_{\mathbf{x}(0)=0}^{\mathbf{x}(\beta)=0} D[\mathbf{x}(\tau)] e^{-S_T} = W[\mathbf{j} = 0, \mathbf{x} - \mathbf{x}' = 0] = W_0, \quad (51)$$

and the first order contribution terms are defined as

$$E_1 = -\frac{1}{\beta W_0} \Sigma_1^0, \quad E_2 = -\frac{1}{\beta W_0} \Sigma_2^0, \quad (52)$$

where $\Sigma_{1,2}^0$ is defined by Eq. (41). While the terms E_1 and E_2 can be evaluated directly using the generating functional, the first term E_0 can not be calculated explicitly from the formula (51). In order to find E_0 one has to note that

$$\int_0^\beta d\tau \int_0^\beta d\tau' Q(\tau - \tau') \langle (\mathbf{x}(\tau) - \mathbf{x}(\tau'))^2 \rangle_0 = -m \frac{d}{dm} e^{-\beta E_0} \quad (53)$$

Using the result (48) we can get following equation for zero order energy

$$\frac{dE_0}{dm} = \frac{3}{4} \frac{\omega^2}{M\Omega}, \quad E_0(m=0) = 0. \quad (54)$$

The solution of the above equation is

$$E_0 = \frac{3}{2}(\Omega - \omega) , \quad E_0 + E_1 = \frac{3}{4} \frac{(\Omega - \omega)^2}{\Omega} . \quad (55)$$

Now it is left to calculate the contribution E_2 . With the help of the formula (47), in the zero temperature limit one can give the following expression for E_2

$$E_2 = - \int_0^\infty d\tau \sum_{\mathbf{k}} \gamma_{\mathbf{k}}^2 \exp \left[-\epsilon(\mathbf{k})\tau - \frac{\mathbf{k}^2}{2} \left(\frac{\omega^2}{\Omega^2} \frac{\tau}{M} + \frac{\Omega^2 - \omega^2}{\Omega^2} \frac{1 - e^{-\Omega\tau}}{M\Omega} \right) \right] \quad (56)$$

First, let us consider the weak coupling limit, i.e. $Q_0 \rightarrow 0$ or $m \rightarrow 0$ so that $\Omega \sim \omega$. Thus we have in the lowest order $E_0 + E_1 = 0$ and the energy E is given by the first expansion term of the ordinary perturbation theory

$$E^{weak} = E_2^{weak} = - \sum_{\mathbf{k}} \gamma_{\mathbf{k}} \frac{1}{\epsilon(\mathbf{k}) + \frac{k^2}{2M}} . \quad (57)$$

In order to prevent the ultraviolet divergence in the above sum over momenta one has to renormalize the coupling constant g according to the second order Born approximation. For this purpose we must add the zero order energy term gn from the Hamiltonian (2) to the energy expression and expand the constant g in powers of scattering length a for the impurity-BEC interaction

$$g = \frac{2\pi a}{m_r} \left(1 + \frac{2a}{\pi} \int dk \right) , \quad m_r = \frac{m_B M}{m_B + M} , \quad (58)$$

The weak coupling energy reexpanded in powers of scattering length is now finite and reads

$$E^{weak} = m_B c^2 \alpha (1 + z) \int_0^\infty dl \left[1 - \frac{1 + z}{\sqrt{1 + 4/l^2} \left(\sqrt{1 + 4/l^2} + z \right)} \right] \quad (59)$$

Here we have introduced dimensionless coupling parameter $\alpha = 4an/(m_B c)$ and the mass ratio $z = m_B/M$. Below we will always use the renormalized expression for E_2 which is

$$E_2^{ren} = E_2 + \frac{4a^2 n}{m_r} \int dk \quad (60)$$

Finally, the whole expression for the ground state energy can be written in the integral form

$$E^{ren}(\Omega, \omega) = \frac{3}{4} \frac{(\Omega - \omega)^2}{\Omega} + \alpha(1 + z) \int_0^\infty dl \left[1 - \frac{1 + z}{2} \frac{l^2}{\sqrt{1 + 4/l^2}} \int_0^\infty ds \exp \left(-\frac{l^2 \sigma_1(s)}{2} \right) \right] , \quad (61)$$

where

$$\sigma_1(s) = s\sqrt{1 + 4/l^2} + sz\frac{\omega^2}{\Omega^2} + \frac{z}{\Omega}\frac{\Omega^2 - \omega^2}{\Omega^2}(1 - e^{-\Omega s}) , \quad (62)$$

The integration in (61) is performed over dimensionless variables s and l , and the energy $E(\Omega, \omega)$ as well as the frequencies Ω and ω are now measured in the units $m_B c^2$.

Following the principal of minimal sensitivity [16] (or Feynman-Jensen variational inequality in case of the energy) we seek the extremum point of the energy with respect to the variational parameters Ω and ω . Fig. (1) shows the energy as a function of Ω , ω for the mass ratio $z = 4/3$, which corresponds He₃ impurity in He₄ BEC, and for the value of coupling constant $\alpha = 5$. This dependence has the form of gutter having a minimum with respect to Ω and not having it with respect to ω . As it can be seen from the Eq. (61), the strong coupling asymptotic of the energy does not depend on ω and behaves like $E_{strong} \sim -\alpha^2$ requiring the optimization with respect to Ω only. On the other hand, the weak coupling regime is independent of both variational parameters. Thus only the intermediate coupling region is sensitive to the choice of the second parameter ω . On Fig. (2) we plotted the energy as a function of the coupling constant α optimized with respect to Ω for different values of the second variational parameter ω . It turns out that at some value of the coupling parameter α the energy becomes negative indicating the existence of the bound impurity/BEC state in spite of the repulsive interaction potential. One can see that the optimized value of the energy rapidly converges to some minimum with increasing ω becoming insensitive to ω at the values of $\omega \geq 50$. Thus the use of this value is safe within variational treatment. So the critical value of the coupling constant α_c defined as $E(\alpha_c) = 0$ decreases from $\alpha_c \sim 3.2$ at $\omega \sim 1$ down to reliable value $\alpha_c \sim 1.8$ at high ω . Next, Fig. (3) shows the critical value of the coupling constant α_c as a function of the mass ratio m_B/M . As one would expect the impurity never self-localizes if its mass is much bigger than the one of the Bose particle and binds with BEC in the weak coupling regime in the opposite case of the small impurity mass.

VI. IMPURITY CORRELATION FUNCTION

Now let us directly consider the correlation function $C(x)$. Using expression for the function Σ_1 , Eq. (48), one can rewrite the expansion (38) up to first cumulant as

$$C_T(\mathbf{x}, \beta \rightarrow \infty) = \exp \left[\beta E_2 - \frac{M\Omega}{8} \left(\frac{\Omega^2 - \omega^2}{\Omega^2} \right)^2 \mathbf{x}^2 + \frac{\Sigma_2(\mathbf{x})}{W_0(\mathbf{x})} \right]. \quad (63)$$

The above function is normalized in such a way that $C_T(\mathbf{x} = 0) = 1$. Then, with the help of the Eqs. (45, 47) the correlation function can be written down in the integral form, namely

$$C_T(\mathbf{x}, \beta \rightarrow \infty) = \exp \left\{ -\frac{z\Omega}{8} \left(\frac{\Omega^2 - \omega^2}{\Omega^2} \right)^2 x^2 + \alpha \frac{(1+z)^2}{2} \int_0^\beta ds \int_0^s ds' \frac{l^2}{\sqrt{1+4/l^2}} \right. \\ \left. \times \exp \left(-\frac{l^2 \sigma_1(s-s')}{2} \right) \left(\frac{\sin(lx\sigma_2(s-s'))}{lx\sigma_2(s-s')} - 1 \right) \right\}, \quad (64)$$

where the function $\sigma_1(s-s')$ is defined by the Eq. (62) and

$$\sigma_2(s-s') = \frac{\omega^2}{\Omega^2} \frac{s-s'}{\beta'} + \frac{1}{2} \frac{\Omega^2 - \omega^2}{\Omega^2} \frac{1}{\sinh \Omega \beta} \\ \times (\sinh \Omega s - \sinh \Omega(\beta - s) - \sinh \Omega s' + \sinh \Omega(\beta - s')) . \quad (65)$$

As in Eq. (61), the integration on the right hand side of the above expression is performed over dimensionless variables s , s' , and l , temperature $1/\beta$ and frequencies Ω and ω are measured in units $m_B c^2$ while the variable x is defined in units $1/m_B c$. Finally, in order to find the best approximation for the correlation function the expansion (64) has to be optimized with respect to the pair of the variational parameters Ω , ω . In general, if the optimization procedure is directly applied to the expansion for the correlation function, the optimized values of variational parameters should depend on the coordinate $\mathbf{x}$. However, it turns out that instead of the variational optimization of the coordinate-dependent correlation function one can use the optimal values of the variational parameters obtained from the optimization of energy. These coordinate-independent values then has to be substituted into the expansion for the correlation function giving the variationally improved result, i.e. convergent strong coupling expansion for the correlation function. This scheme of the variational perturbation theory was successfully applied for the variational solutions of time-dependent equations in non-linear dynamics [17, 18].

We will also define the mean value of x^2 representing the square of the radius of localization or polaron radius as

$$\langle x^2 \rangle = \frac{\int x^2 C(\mathbf{x}) d\mathbf{x}}{\int C(\mathbf{x}) d\mathbf{x}} \quad (66)$$

Fig. (4) shows the mean square of the radius of localization as a function of coupling constant. It has inverse dependence on coupling strength. The inset shows the correlation function itself for different values of coupling strength. The correlation function has Gaussian-like shape regardless of α which means that formally the particle is always localized. This is the consequence of the choice of the trial action used in this method. However, this correlation function should converge to the true-one at large values of interaction strength. It is interesting to examine the localization radius at the coupling constant α_e at which the energy has maximum, i.e. $dE(\alpha)/d\alpha|_{\alpha_e} = 0$. We notice that $\langle x^2(\alpha_e) \rangle \sim 1/m_{BC}$. This can be expressed through a so called healing length of the BEC $\xi = 1/zp_c$, where p_c is the critical momentum of the impurity above which the dissipation takes place in case of real-time dynamics in accordance to Landau's criterion. This leads to $\sqrt{\langle x^2(\alpha_e) \rangle} \sim 1/p_c$, or using Heisenberg uncertainty for the localized particle $\sqrt{\langle p^2 \rangle} \sim p_c$, i.e. the average momentum of a particle at the point of self localization is of the order of the critical momentum. One can make physical sense of this fact by noticing that this means that the tendency of self-localization appears when the exchange of energy between particle and BEC becomes possible.

VII. CONCLUSION

In this work we consider a problem of self-localization of impurity in degenerate BEC. This problem has been considered previously using the mean-field approach. Here we use a full quantum description of the ground state of impurity surrounded by BEC. The variational perturbation method is employed to calculate the imaginary-time propagator of impurity in degenerate BEC. The free energy and the spatial correlation of impurity in BEC is obtained as a function of coupling strength. Our results point to a possible relation between self localization and the real-time particle/BEC energy exchange. In this work we explicitly used the degeneracy of the BEC. However, the methodology developed in this paper can be extended to Bose systems without an assumption of diluteness. The latter can be done by using the four-point Green's function of the strong coupled Bose liquid [19] in the leading expansion term instead of degenerate BEC propagator used in this work.

VIII. ACKNOWLEDGMENTS

This work has been supported by the NSF CAREER award ID 0645340.

- [1] A. P. Chikkatur, A. Görlitz, D. M. Stamper-Kurn, S. Inouye, S. Gupta and W. Ketterle, Phys. Rev. Lett. **85** 483 (2000)
- [2] S. Grebenev, P. Toennis and A. Vilesov, Science **279** 2083 (1998)
- [3] S. Grebenev, B. Sartakov, P. Toennis and A. Vilesov, Science **289** 1532 (2000)
- [4] R. Kalas and D. Blume, Phys. Rev. A **73** 043608 (2006)
- [5] F. M. Cucchietti and E. Timmermans, Phys. Rev. Lett. **96** 210401 (2006)
- [6] K. Sacha and E. Timmermans, Phys. Rev. A **73** 063604 (2006)
- [7] M. Bruderer, W. Bao and D. Jaksch, arXiv: 0801.4000v4 (2008)
- [8] R. Feynman, Phys. Rev. **97** 660 (1955)
- [9] V. Sa-yakanit, V. Lakhno and K. Hass Phys. Rev. B **57** 5772 (1998)
- [10] R. Rosenfelder and A. Schreiber, arXiv: ntl-th/9504002v1 (1995)
- [11] J. Devreese, in *Fluctuating Paths and Fields*, 3rd ed., (World Scientific, Singapore, 2001)
- [12] N. N. Bogoliubov, J. Phys. (Moscow) **11** 23 (1947)
- [13] A. Abrikosov, L. Gorkov and I. Dzyaloshinski, *Methods of Quantum Field Theory in Statistical Physics*, (Dover Publication, New York, 1975)
- [14] A. Miller, D. Pines and P. Nozieres, Phys. Rev. **127** 1452 (1962)
- [15] A. Novikov and M. Ovchinnikov, J. Phys. A: Math. Theor. **42** 135301 (2009)
- [16] H. Kleinert, *Path Integrals in Quantum Mechanics, Statistics, Polymer Physics and Financial Markets*, 3rd ed., (World Scientific, 2004)
- [17] S. Brandt, A. Pelster and R. Wessel, Phys. Rev. E **74** 036201 (2006)
- [18] A. Pelster, A. Novikov, U. Kleinekathöfer and M. Schreiber (unpublished results, 2005)
- [19] N. Bogoliubov, C. Malyshev, R. Bullough and J. Timonen, arXiv: cond-mat/0306735v1 (2003)

Figures




FIG. 1: Energy as a function of variational parameters Ω and ω as described by Eq. (61). $\text{Alpha}=1$, $z=4/3$




FIG. 2: Energy optimized with respect to Ω as a function of the coupling constant α for different values of second variational parameter ω .




FIG. 3: Critical value of coupling constant α_c as a function of the mass ratio m_B/M .




FIG. 4: Mean value of x^2 as a function of coupling constant, x is shown in units $1/m_B c$. Inset - correlation function $c(x)$ for different values of coupling constant $\alpha = 0.5$ (upper curve), $\alpha = 1, 5$ and $\alpha = 10$ (lower curve).